

# Microsurgical Resection of Suprasellar Craniopharyngioma by Pterional Approach: 3-Dimensional Operative Video

Ricardo Lourenço Caramanti, MD, MSc<sup>\*,†</sup>, Erica Antunes Effgen, MD<sup>\*</sup>, Raysa Moreira Aprigio, MD<sup>\*</sup>, Dionei Freitas de Moraes, MD, PhD<sup>\*</sup>, Carlos Eduardo Dall Aglio Rocha, MD<sup>\*</sup>, Eberval Gadelha Figueiredo, MD, PhD<sup>‡</sup>, Feres Chaddad-Netto, MD, PhD<sup>§</sup>

<sup>\*</sup>Department of Neurosurgery, São José do Rio Preto Medical School (FAMERP), São José do Rio Preto, São Paulo, Brazil; <sup>†</sup>Department of Neurosurgery, Universidade de São Paulo (USP), São Paulo, Brazil; <sup>‡</sup>Department of Neurosurgery, Universidade Federal de São Paulo (UNIFESP), São Paulo, Brazil

**Correspondence:** Ricardo Lourenço Caramanti, MD, MSc, Department of Neurosurgery, São José do Rio Preto Medical School, Brigadeiro Faria Lima Av. 5544, São José do Rio Preto, São Paulo, 15090-000, Brazil. Email: ricardocaramanti1987@gmail.com

**Received,** December 30, 2022; **Accepted,** March 15, 2023; **Published Online,** May 24, 2023.

© Congress of Neurological Surgeons 2023. All rights reserved.

Craniopharyngioma is an uncommon slow-growing lesion that comprised 2% of all intracranial tumors. Approximately 60% of cases are diagnosed after 16 years despite bimodal age distribution between 5 to 15 and 45 to 60 years. Typical locations are suprasellar in 75% of cases, 21% sellar and suprasellar, rarely restricted to sella.

There are 2 histological types: adamantinomatous, which most commonly appears in children, and papillary, more common in adults.

Clinical symptoms are associated with structures such as pituitary, third ventricle, optic, and oculomotor nerve compression by mass effect, generating hormonal disturbances, hydrocephalus, visual deficits, and raised intracranial pressure signs.

Tumor adhesion to Willis polygon and optic nerves can be challenging, increasing bleeding, ischemia, and loss of vision risks. We present a 35-year-old patient with a history of frontal headache associated with 5 months of amenorrhea and a mouth of progressive visual loss. Physical examination showed bitemporal hemianopsia. The neurological investigation by magnetic resonance imaging revealed a sellar/suprasellar solid cystic tumor with pituitary compression, without third ventricle invasion, which was compatible with craniopharyngioma. The patient underwent resection through a pterional approach, and the lesion was resected entirely without symptoms worsening. A pathology examination confirms craniopharyngioma diagnosis.

In this 3-dimensional video, the authors present the step-by-step microsurgical technique to perform a sellar/suprasellar craniopharyngioma resection by pterional approach.

The patient consented to the procedure and publication of his images.

**Patient consent:** The patient signed the institutional consent form, which allows the use of his/her images and videos for any medical publications in conferences and/or scientific.

**KEY WORDS:** Craniopharyngioma, Pterional approach, Interpeduncular cistern, Hemianopsia

*Operative Neurosurgery* 25:E155–E156, 2023

<https://doi.org/10.1227/ons.0000000000000756>

Watch now at <http://dx.doi.org/10.1227/ons.0000000000000756>

## Funding

This study did not receive any funding or financial support.

## Disclosures

The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

## COMMENTS

The authors present a case of a patient with a craniopharyngioma who underwent a pterional approach for resection of the tumor. The surgery is beautifully done and highlights some important points. Given that these tumours can vary quite a bit from patient to patient, it is important to understand the relationship of the tumor to the critical structures